



Rituximab, bendamustine, and low-dose cytarabine as induction therapy in elderly patients with mantle cell lymphoma: a multicentre, phase 2 trial from Fondazione Italiana Linfomi.

Lancet Haematol. 2017 Jan;4(1):e15-e23. doi: 10.1016/S2352-3026(16)30185-5. Epub 2016 Dec 5.

Published: 13 July 2017 -PMID: 27927586

Visco C¹, Chiappella A², Nassi L³, Patti C⁴, Ferrero S⁵, Barbero D⁵, Evangelista A⁶, Spina M⁷, Molinari A⁸, Rigacci L⁹, Tani M¹⁰, Rocco AD¹¹, Pinotti G¹², Fabbri A¹³, Zambello R¹⁴, Finotto S¹⁵, Gotti M¹⁶, Carella AM¹⁷, Salvi F¹⁸, Pileri SA¹⁹, **Ladetto M¹⁸**, Ciccone G⁶, Gaidano G³, Ruggeri M²⁰, Martelli M¹¹, Vitolo U².

Author information:

1. Department of Cell Therapy and Hematology, San Bortolo Hospital, Vicenza, Italy. Electronic address: carlovisco@hotmail.com.
2. Hematology, Citta' della salute e della scienza University Hospital, Torino, Italy.
3. Hematology, Department of Translational Medicine, University of Eastern Piedmont, Novara, Italy.
4. Hematology, Ospedali Riuniti Villa Sofia-Cervello, Palermo, Italy.
5. Molecular Biotechnologies and Health Sciences, University of Torino, Torino, Italy.
6. Clinical Epidemiology, Città della Salute e della Scienza and CPO Piemonte, Torino, Italy.
7. Medical Oncology A, National Cancer Institute, Aviano, Italy.
8. Hematology, Ospedale degli Infermi, Rimini, Italy.
9. Hematology, University of Florence, Florence, Italy.
10. Hematology, Ospedale di Ravenna, Ravenna, Italy.
11. Cellular Biotechnologies and Hematology, Sapienza University, Rome, Italy.
12. Oncology ASST-settelaghi, Varese, Italy.
13. Hematology, S Maria alle Scotte University Hospital, Siena, Italy.
14. Hematology, University of Padova, Padova, Italy.
15. Medical Oncology 1, Veneto Institute of Oncology-IRCCS, Padova, Italy.
16. Hematology Oncology, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy.
17. past-director hematology unit IRCCS San Martino, Genova, Italy.
18. **Hematology, SS Antonio e Biagio e Cesare Arrigo Hospital, Alessandria, Italy.**
19. Haematopathology Unit, European Institute of Oncology, Milan, Italy; Bologna University School of Medicine, Bologna, Italy.
20. Department of Cell Therapy and Hematology, San Bortolo Hospital, Vicenza, Italy.

Abstract

BACKGROUND:

The combination of rituximab, bendamustine, and cytarabine (R-BAC) was highly active in a pilot trial of mantle cell lymphoma, but its use was restricted by high haematological toxicity. We aimed to assess the efficacy and safety of an R-BAC regimen with low-dose cytarabine (RBAC500).

METHODS:

In this multicentre, phase 2 trial, we recruited previously untreated patients with an established histological diagnosis of mantle cell lymphoma from 29 Fondazione Italiana Linfomi centres in Italy. Patients had to be older than 65 years and fit according to the comprehensive geriatric assessment, or aged 60-65 years if they were ineligible for high-dose chemotherapy plus autologous stem-cell transplantation and were fit or unfit. All patients received RBAC500 (rituximab 375 mg/m² on day 1, bendamustine 70 mg/m² on days 2 and 3, and cytarabine 500 mg/m² on days 2-4; all administered intravenously) every 4 weeks for up to six cycles. Primary endpoints were the proportion of patients achieving complete response at the end of treatment and toxicity, defined as the occurrence of any of the stop treatment criteria or of any episode of relevant toxicity. All patients who started at least one cycle of RBAC500 were included in the primary and safety analyses. Using efficacy and toxicity as a composite primary endpoint, we considered the final conclusion positive if more than 28 of 57 patients achieve complete response and fewer than 18 of 57 patients report toxicities. This study is registered with EudraCT, number 2011-005739-23, and ClinicalTrials.gov, number [NCT01662050](https://clinicaltrials.gov/ct2/show/study/NCT01662050), and is completed.

FINDINGS:

Between May 2, 2012, and Feb 25, 2014, we enrolled 57 patients (median age 71 years, IQR 67-75). 54 (95%) patients received at least four RBAC500 cycles (three discontinued because of toxicity), and 38 (67%) completed six cycles. Two (4%) had disease progression (one after the fourth cycle and one after the sixth cycle). All 52 (91%, lower limit of one-sided 95% CI 85%) remaining patients achieved complete response at the end of treatment. 23 (40%, upper limit of one-



sided 95% CI 53%) of 57 patients had at least one episode of relevant toxicity. The most frequent grade 3-4 haematological toxicities were neutropenia (149 [49%] of 304 cycles) and thrombocytopenia (158 [52%]). Most treatment-related non-haematological adverse events were of grade 1-2, with the most frequent ones being fatigue (14 [25%] patients), nausea or vomiting (12 [21%]), and infusion-related reactions or tumour lysis syndrome (12 [21%]). 41 (72%) patients required a dose reduction. 12 patients died during the study, but no deaths were related to treatment.

INTERPRETATION:

RBAC500 is an effective treatment for elderly patients with mantle cell lymphoma and, despite not meeting our prespecified safety boundary, haematological toxicity was manageable with appropriate supportive care and dose reduction. Since maintenance therapy is not required, RBAC500 could be considered an option and should be studied in phase 3 trials.

FUNDING:

Fondazione Italiana Linfomi and Mundipharma.